

Chen, Z., Xu, Xianfa, Zhang, J., Liu, Y., Xu, Xianggang, et al. 2016. Application of LC-MS-based global metabolomic profiling methods to human mental fatigue. *Analytical Chemistry*. <https://doi.org/10.1021/acs.analchem.6b03421>.

Dresch, V., Sánchez-López, M. del P., & Saavedra, A. I. 2015. Occupational Stress versus Cortisol Associated with Psychological Variables and Gender Differences among Women and Men in Nursing. *Health*, 07(07): 838-846.

Du, C.-L., Lin, M. C., Lu, L., & Tai, J. J. 2011. Correlation of Occupational Stress Index with 24-hour Urine Cortisol and Serum DHEA Sulfate among City Bus Drivers: A Cross-sectional Study. *Safety and Health at Work*, 2(2): 169-75.

Eck, M. van, Berkhof, H., Nicolson, N., & Sulon, J. 1996. The Effects of Perceived Stress, Traits, Mood States, and Stressful Daily Events on Salivary Cortisol. *Psychosomatic Medicine*, 58(5): 447-458.

Limm, H., Gundel, H., Heinmuller, M., Marten-Mittag, B., Nater, U. M., et al. 2010. Stress management interventions in the workplace improve stress reactivity: a randomised controlled trial. *Occupational and Environmental Medicine*, 68(2): 126-133.

Limm, H., Gündel, H., Heinmüller, M., Marten-Mittag, B., Nater, U. M., et al. 2011. Stress management interventions in the workplace improve stress reactivity: a randomised controlled trial. *Occupational and Environmental Medicine*, 68(2): 126.

Minhee, S. 2017. An Exploratory Study on Occupational Stress and Anxiety Through Salivary Cortisol and Self-Report Scale in Korean Nurses on Shift and Regular Work. *Journal of Korean Biological Nursing Science*, 19(3): 206-213.

Moustaka, E., Maria, M., & Constantinidis, T. 2015. Measuring Occupational Stress and HRA Axis Dysregulation among Healthy Workers by Salivary Cortisol Levels. *British Journal of Medicine and Medical Research*, 6(11): 1040-1051.

Reale, M., Costantini, E., D'Angelo, C., Coppeta, L., Mangifesta, R., et al. 2020. Network between Cytokines, Cortisol and Occupational Stress in Gas and Oilfield Workers. *International Journal of Molecular Sciences*, 21(3): 1118.

Rocha, M. C. P. da, Martino, M. M. F. D., Grassi-Kassisse, D. M., & Souza, A. L. de. 2013. Stress among nurses: an examination of salivary cortisol levels on work and day off. *Revista Da Escola de Enfermagem Da USP*, 47(5): 1187-1194.

Rotter, M., Brandmaier, S., Covic, M., Burek, K., Hertel, J., et al. 2018. Night Shift Work Affects Urine Metabolite Profiles of Nurses with Early Chronotype. *Metabolites*, 8(3): 45.

Sood, P., Priyadarshini, S., & Aich, P. 2013. Estimation of Psychological Stress in Humans: A Combination of Theory and Practice. *PLoS ONE*, 8(5). <https://doi.org/10.1371/journal.pone.0063044>.

Susoliakova, O., Smejkalova, J., Bicikova, M., Hodacova, L., Malkova, A., et al. 2018. Assessment of work-related stress by using salivary cortisol level examination among early morning shift workers. *Central European Journal of Public Health*, 26(2): 92-97.

Walvekar, S. S., Ambekar, J. G., & Devaranavadagi, B. B. 2015. Study on serum cortisol and perceived stress scale in the police constables. *Journal of Clinical and Diagnostic Research: JCDR*, 9(2): BC10-4.

Wirtz, P. H., Ehlert, U., Kottwitz, M. U., Marca, R. L., & Semmer, N. K. 2013. Occupational role stress is associated with higher cortisol reactivity to acute stress. *Journal of Occupational Health Psychology*, 18(2): 121-31.